arXiv:1512.02681v1 [math.GR] 8 Dec 2015

# NEGATIVE DEFINITE FUNCTIONS ON GROUPS WITH POLYNOMIAL GROWTH

FABIO CIPRIANI. AND JEAN-LUC SAUVAGEOT

Abstract. The aim of this work is to show that on a locally compact, second countable, compactly generated group $G$ with polynomial growth and homogeneous dimension $d_h$, there exist a continuous, proper, negative definite function $\ell$ with polynomial growth dimension $d_\ell$ arbitrary close to $d_h$.

## 1. Introduction and statement of the results

The length function $\ell$ associated to a finite set of generators of a countable, discrete, finitely generated group $\Gamma$, may be used to reveal interesting aspects of the group itself. This is the spirit of metric group theory (see [12]). In a famous result of U. Haagerup [11], for example, the length functions on the free groups $\mathbb{F}_n$ with $n \geq 2$ generators, has been used to prove that, even if these groups are not amenable, they still have an approximation property: there exists a sequence $\varphi_n \in c_0(\mathbb{F}_n)$ of normalized, positive definite functions, converging pointwise to the constant function1. Correspondingly, the trivial representation is weakly contained in a $C_0$-representation (one whose coefficients vanish at infinity) (see [10]). These properties of $\mathbb{F}_n$ only depend to the fact that the length function is *negative definite and proper*. The class of groups where a length function has the latter properties has been characterized in several independent ways and form the object of intensive investigations (see [2]). On another, somewhat opposite, side, a deeply studied class of groups are those possessing the Kazhdan property (T), i.e. those where *every negative definite function is bounded* (see for example [1]). Finer asymptotic properties of length functions are connected to other fundamental properties of groups. For example, if $\ell$ has *polynomial growth* (see below for the definition) then $\Gamma$ is *amenable* and its von Neumann algebra $vN(\Gamma)$ is *hyperfinite* (weak closure on an increasing sequence of matrix algebras).

However, length functions are not necessarily negative definite and one may wonder how far is a given length function from being negative definite.

Reversing the point of view, in the present work we prove that on finitely generated, discrete groups $\Gamma$ or, more generally, on compactly generated, locally compact, second countable groups $G$, with polynomial growth, there always exists a continuous, negative definite function with a (polynomial) growth arbitrarily close to the one of $\Gamma$ or $G$.

The consideration of negative definite functions and their growth properties are also meaningful in Noncommutative Potential Theory (see [3], [6]) and Quantum Probability (see [4]). As an application of the main results, we provide a natural construction of a noncommutative Dirichlet form on the group von Neumann algebra $\lambda(G)''$ with an upper spectral growth dimension $d_S$ bounded above by the polynomial growth homogeneous dimension $d_h$ of $G$.

*Date*: December 8, 2015.

1991 *Mathematics Subject Classification.* Primary 20F65; Secondary 20F69, 57M07.

*Key words and phrases.* Group, polynomial growth, negative definite function, homogeneous dimension.

This work was supported by Italy I.N.D.A.M. France C.N.R.S. G.D.R.E.-G.R.E.F.I. Geometrie Noncommutative and by Italy M.I.U.R.-P.R.I.N. project N. 2012TC7588-003.

Let $G$ be a locally compact, second countable, compactly generated group with identity $e \in G$ and let $\mu$ be one of its left Haar measures.

The main results of the work are the following.

**Theorem 1.1.** *Suppose that $G$ has polynomial growth and homogeneous dimension $d_h$. Then, for all $d > d_h$, there exists a continuous, proper, negative type function $\ell$ with polynomial growth such that*

$$\mu\{s \in G \,|\, \ell(s) \le x\} = O(x^d)\,, \quad x \to +\infty. \tag{1.1}$$

**Theorem 1.2.** *Suppose that $G$ has polynomial growth and homogeneous dimension $d_h$. Then, there exists a continuous, proper, negative type function $\ell$ with polynomial growth such that*

$$\forall d > d_h\,, \quad \mu\{s \in G \,|\, \ell(s) \le x\} = O(x^d) \qquad x \to +\infty. \tag{1.2}$$

As a straightforward application of these results to Noncommutative Potential Theory we have the following corollary. Further investigations will be discussed in [7].

**Corollary 1.3.** *Let $\Gamma$ be a discrete, finitely generated group with polynomial growth and homogeneous dimension $d_h$. Let $\lambda(\Gamma)''$ be the von Neumann algebra generated by the left regular representation and let $\tau$ be its trace.*
*Then there exists on $L^2(\lambda(\Gamma)'', \tau)$ a noncommutative Dirichlet form $(\mathcal{E}, \mathcal{F})$ with discrete spectrum and upper spectral dimension $d_S \le d_h$.*

Here the *upper spectral dimension* $d_S$ of the Dirichlet form is defined as

$$d_S := \inf\{d > 0 : \limsup_{x \to +\infty} \sharp\{\lambda \in \mathrm{sp}\,(L) : \lambda \le x\}/x^d < +\infty\}$$

where $\mathrm{sp}(L)$ denotes the spectrum of the self-adjoint, nonnegative operator $(L, \mathrm{dom}(L))$ associated to $(\mathcal{E}, \mathcal{F})$

$$\mathcal{E}[a] = \|\sqrt{L}a\|_2^2 \qquad a \in \mathcal{F} = \mathrm{dom}(L)\,.$$

## 2. Groups with polynomial growth

Let $K \subset G$ be a compact, symmetric, generator set with non empty interior

$$G = \bigcup_{n=1}^{\infty} K^n\,, \quad K^{-1} = K\,, \quad K^\circ \neq \emptyset\,,$$

where

$$K^{-1} := \{s^{-1} \in G : s \in K\}\,, \qquad K^n := \{s_1 \dots s_n \in G : s_k \in K\,, k = 1, \dots, n\}\,.$$

We may always assume that $K$ is a neighborhood of the identity $e \in K^\circ \subset G$.
Let us recall that $G$ has *polynomial growth* if

$$\exists\, c, d > 0 \quad \text{such, that} \quad \mu(K^n) \le c\,(n+1)^d\,, \qquad n \ge 1 \tag{2.1}$$

or, equivalently, if

$$\frac{1}{\log(n+1)} \log(\mu(K^n)) \le d + \frac{1}{\log(n+1)} \log(c)\,, \qquad n \ge 1 \tag{2.2}$$

and in that case its *homogeneous dimension* is defined as

$$d_h = \limsup_{n \to \infty} \frac{1}{\log n} \log(\mu(K^n))\,. \tag{2.3}$$

The class of locally compact, second countable, compactly generated groups with polynomial growth includes nilpotent, connected, real Lie groups and finitely generated, nilpotent, countable discrete groups (see [8] and [12] Chapter VII $\mathcal{S}$ 26). For example, the homogeneous dimension of the discrete Heisenberg group is 4 (see [12] Chapter VII $\mathcal{S}$ 21).

Recall also the M. Gromov's characterization [9] of finitely generated, countable discrete groups with polynomial growth as those which have a nilpotent subgroup of finite index (see also [12] Chapter VII $\mathcal{S}$ 29 and [13] for a generalization to locally compact groups).

*Remark* 2.1. Let us observe that since $K$ is a neighborhood of the identity $e \in G$, we have $K^{n-1} \subset K^{n-1}K^\circ \subset (K^n)^\circ$ so that

$$\bigcup_{k=0}^{\infty} (K^n)^\circ = G$$

and we have an open cover of $G$. As a consequence

*every compact set in $G$ is contained in $K^n$ for some $n \geq 1$.*

## 3. Proof of the Theorems

In the following, we will consider the sequence $\{\alpha_n\}_{n=1}^\infty \subset [0, +\infty)$ defined by

$$\alpha_1 := 0\,, \qquad \frac{\mu(K^n)}{\mu(K^{n-1})} =: 1 + \alpha_n\,, \qquad n \geq 2\,.$$

Equation (2.2) can thus be written as

$$\frac{1}{\log(n+1)} \sum_{k=1}^{n} \log(1+\alpha_k) \leq d + \frac{1}{\log(n+1)} \log(c) \leq d'\,, \qquad n \geq 1\,, \tag{3.1}$$

with $d' := d + \log_2 c$.

3.1. **Some preliminary estimates.** The proof of the theorems relies on the following lemmas.

**Lemma 3.1.** *For any $\beta \in (0,1)$ the set*

$$E_\beta := \left\{ n \in \mathbb{N}^* \,:\, \alpha_n > n^{-\beta} \right\}$$

*has vanishing density.*

*Proof.* Let us fix $K > 0$ such that $\log(1 + n^{-\beta}) \geq K n^{-\beta}$ for all $n \geq 1$. By equation (3.1), we have

$$d' \log(n+1) \geq \sum_{k \in [1,n] \cap E_\beta} \log(1+\alpha_k) \geq K n^{-\beta} \sharp\{[1,n] \cap E_\beta\}$$

so that

$$\frac{1}{n} \sharp\{[1,n] \cap E_\beta\} \leq \frac{d'}{K} \frac{\log(n+1)}{n^{1-\beta}} \to 0 \quad \text{as} \quad n \to \infty\,.$$

□

**Lemma 3.2.** *For any $\beta \in (0,1)$ and $\gamma > 1/\beta$, the set*

$$F_{\beta,\gamma} := \left\{ n \in \mathbb{N} \,:\, [\, n^\gamma, (n+1)^\gamma\,] \cap \mathbb{N} \subset E_\beta \right\}$$

*has vanishing density.*

*Proof.* For any $\varepsilon \in (0,1/2)$ and any integer $N \geq 2$, we have

$$\begin{aligned} d' &\geq \frac{1}{\log(N+1)^\gamma} \sum_{(\varepsilon N)^\gamma \leq k \leq N^\gamma} \log(1+\alpha_k) \\ &\geq \frac{1}{\log(N+1)^\gamma} \sum_{n \in [\varepsilon N, N-1] \cap F_{\beta,\gamma}} \sum_{k \in [n^\gamma, (n+1)^\gamma)} \log(1+\alpha_k) \\ &\geq \frac{K}{\gamma \log(N+1)} \sum_{n \in [\varepsilon N, N-1] \cap F_{\beta,\gamma}} \sum_{k \in [n^\gamma, (n+1)^\gamma)} k^{-\beta} \qquad N \geq 1\,. \end{aligned}$$

For any fixed $n \in [\varepsilon N, N-1] \cap F_{\beta,\gamma}$, there exists $t \in (n, n+1)$ such that $(n+1)^\gamma - n^\gamma = \gamma t^{\gamma-1}$. As $n \geq \varepsilon N$, we have $t/N > n/N \geq \varepsilon$ so that for $n \in [\varepsilon N, N-1] \cap F_{\beta,\gamma}$ the following bound holds true

$$\begin{aligned} \sum_{n^\gamma \leq k < (n+1)^\gamma} k^{-\beta} &\geq N^{-\beta\gamma}\big((n+1)^\gamma - n^\gamma - 2\big) \\ &= N^{-\beta\gamma}\big(\gamma t^{\gamma-1} - 2\big) \\ &> N^{-\beta\gamma}\big(\gamma(\varepsilon N)^{\gamma-1} - 2\big). \end{aligned}$$

Thus we have

$$\begin{aligned} d' &\geq \frac{K}{\gamma \log(N+1)} \sum_{n \in [\varepsilon N, N-1] \cap F_{\beta,\gamma}} \Big(\gamma \varepsilon^{\gamma-1} N^{(1-\beta)\gamma-1} + o(N^{(1-\beta)\gamma-1})\Big) \\ &\geq \frac{K}{\log(N+1)} \Big(\varepsilon^{\gamma-1} N^{(1-\beta)\gamma-1} + o(N^{(1-\beta)\gamma-1})\Big) \sharp\{[\varepsilon N, N-1] \cap F_{\beta,\gamma}\} \end{aligned}$$

and we may deduce

$$\frac{d'}{K\varepsilon^{1-\gamma}} \frac{\log(N+1)}{N^{(1-\beta)\gamma}} \Big(1 + o(1)\Big) \geq \frac{\sharp\{[\varepsilon N, N-1] \cap F_{\beta,\gamma}\}}{N}\,.$$

For $N$ big enough, we have

$$\frac{\sharp\{[\varepsilon N, N-1] \cap F_{\beta,\gamma}\}}{N} \leq \varepsilon$$

and finally

$$\frac{\sharp\{[1, N-1] \cap F_{\beta,\gamma}\}}{N} \leq \frac{\sharp\{[1, \varepsilon N] \cap F_{\beta,\gamma}\}}{N} + \frac{\sharp\{[\varepsilon N, N-1] \cap F_{\beta,\gamma}\}}{N} \leq 2\varepsilon\,.$$

□

3.2. **Proofs of the theorems.** To build up a negative type function having the properties required in Theorem 1.2, we begin to construct a suitable sequence of positive definite functions.

Fix $\beta \in (0,1)$ and $\gamma > 1/\beta$. For any fixed $n \notin F_{\beta,\gamma}$, we can chose $k(n) \in [\, n^\gamma, (n+1)^\gamma\,] \backslash \mathbf{E}_\beta$ in such a way that

$$n^\gamma \leq k(n) \leq (n+1)^\gamma\,, \qquad k(n)^{-\beta} \geq \alpha_{k(n)} = \frac{\mu(K^{k(n)+1}) - \mu(K^{k(n)})}{\mu(K^{k(n)})}\,.$$

Denote by $\mathcal{B}(L^2(G,\mu))$ the Banach algebra of all bounded operators on the Hilbert space $L^2(G,\mu)$ and by $\lambda : G \to \mathcal{B}(L^2(G,\mu))$ the left regular unitary representation of $G$, defined as $(\lambda(s)a)(t) := a(s^{-1}t)$ for $a \in L^2(G,\mu)$ and $s, t \in G$.

Consider now the function $\xi_n := \mu(K^{k(n)})^{-1/2}\chi_{K^{k(n)}} \in L^2(G,\mu)$, with unit norm, and set

$$\omega_n : G \to \mathbb{R} \quad \omega_n(s) :=< \xi_n, \lambda(s)\xi_n >_{L^2(G)} = \frac{\mu(sK^{k(n)} \cap K^{k(n)})}{\mu(K^{k(n)})}\,.$$

By construction we have that
• $\omega_n \in C_c(G)$ is a continuous, normalized, positive definite function with compact support. Moreover
• $\omega_n(s) \geq 1 - n^{-\beta\gamma}$ for all $s \in K$.

In fact, by the translation invariance of the Haar measure and since $K^{k(n)} \supset sK^{k(n)-1}$ for $s \in K$, we have

$$\begin{aligned}\omega_n(s) &= \frac{\mu(sK^{k(n)} \cap K^{k(n)})}{\mu(K^{k(n)})} \geq \frac{\mu(sK^{k(n)} \cap sK^{k(n)-1})}{\mu(K^{k(n)})} = \frac{\mu(s(K^{k(n)} \cap K^{k(n)-1})}{\mu(K^{k(n)})} \\ &= \frac{\mu(K^{k(n)} \cap K^{k(n)-1})}{\mu(K^{k(n)})} = \frac{\mu(K^{k(n)-1})}{\mu(K^{k(n)})} = (1+\alpha_{k(n)})^{-1} \geq 1 - \alpha_{k(n)} \\ &\geq 1 - k(n)^{-\beta} \geq 1 - n^{-\beta\gamma}\,.\end{aligned}$$

**Lemma 3.3.** *For $n \notin F_{\beta,\gamma}$ and any integer $p \geq 1$, we have*

$$\omega_n(s) \geq 1 - pn^{-\beta\gamma} \qquad s \in K^p\,. \tag{3.2}$$

*Proof.* Consider $s = s_1 \cdots s_p \in K^p$ for some $s_1, \cdots, s_p \in K$ and notice that

$$\begin{aligned}2\big(1-\omega(s)\big) &= ||\xi_n - s\xi_n||^2 = ||\xi_n - s_1 \cdots s_p\xi_n||^2 \\ &= ||\xi_n - s_1\xi_n + s_1\xi_n - s_1s_2\xi_n + \cdots + s_1 \cdots s_{p-1}\xi_n - s_1 \cdots s_p\xi_n||^2 \\ &\leq \big(||\xi_n - s_1\xi_n|| + ||\xi_n - s_2\xi_n|| + \cdots + ||\xi_n - s_p\xi_n||\big)^2 \\ &\leq p^2 \sup_{\sigma\in K} ||\xi_n - \sigma\xi_n||^2 \\ &= 2p^2 \sup_{\sigma\in K} \big(1 - \omega_n(\sigma)\big) \leq 2p^2 n^{-2\beta\gamma}.\end{aligned}$$

□

*Proof of Theorem 1.1.* By previous lemma, the series $\sum_{n\notin F_{\beta,\gamma}} \big(1-\omega_n(s)\big)$ converges uniformly on $K^p$ for any integer $p \geq 1$. Hence, it converges uniformly on each compact subset of $G$ and

$$\ell : G \to [0,+\infty) \qquad \ell(s) = \sum_{n\notin F_{\beta,\gamma}} \big(1 - \omega_n(s)\big)$$

is a continuous, negative definite function. The function $\ell$ is proper: in fact, if $m \in \mathbb{N}$ is greater or equal to the integer part of $(N+1)^\gamma$, then, for $s \notin K^{2m}$, we have $\omega_n(s) = 0$ for all $n \in [1,N]\backslash F_{\beta,\gamma}$ so that

$$\ell(s) \geq \sharp\{\,[1,N]\backslash F_{\beta,\gamma}\} \qquad s \notin K^{2m}\,. \tag{3.3}$$

According to Lemma 3.2 we can write $\sharp\{\,[1,N]\backslash F_{\beta,\gamma}\} = N(1-\varepsilon_N)$ for $\varepsilon_N \to 0$. The set $\{s \,|\, \ell(s) \leq N(1-\varepsilon_N)\,\}$ is contained in $K^{2m}$, where $m$ is the integer part of $(N+1)^\gamma + 1$, whose volume $\leq c\,2^d\big(N+1)^\gamma + 1\big)^d = O(N^{\gamma d})$ For fixed $\varepsilon > 0$ and $N$ large enough, we have, for $x \in [N-1, N]$,

$$\mu\{\ell \leq x(1-\varepsilon)^{-1}\} \leq c' N^{\gamma d} \leq c'' x^{\gamma d}$$

so that $\mu\{\ell \leq x\} \leq c'' x^{\gamma d}$. □

*Proof of Theorem 1.2.* Let us chose a decreasing sequence $d_m$ converging to $d_h$, and for all $m$, chose a function $\ell_m$ such that $\ell_m(x) = O(x^{d_m})$, $x \to +\infty$ and $\sup_K \ell_d(x) = 1$ (by normalization). Hence $\ell \le k^2$ on $K^k$. It is enough to set $\ell = \sum_m 2^{-m}\ell_m : \{\ell \le x\} \subset \{\ell_m \le 2^m x\}$, whose measure is $O(x^{d_m})$.

□

*Proof of Corollary 1.3.* Recall first that the GNS space $L^2(\lambda(\Gamma)'', \tau)$ can be identified, at the Hilbert space level, with $l^2(\Gamma)$. Consider now the continuous, proper, negative type function $\ell$ constructed in Theorem 1.2. By [5 $\mathcal{S}$ 10.2], the quadratic form

$$\mathcal{E} : \mathcal{F} \to [0, +\infty) \qquad \mathcal{E}[a] = \sum_{s\in\Gamma} \ell(s)\,|a(s)|^2\,,$$

defined on the subspace $\mathcal{F} \subset l^2(\Gamma)$ where the sum is finite, is a noncommutative Dirichlet form whose associated nonnegative, self-adjoint operator on $l^2(\Gamma)$ is the multiplication operator by $\ell$ given by

$$\mathrm{dom}(L) := \{a \in l^2(\Gamma) : \ell \cdot a \in L^2(G, \mu)\} \quad (La)(s) := \ell(s) \cdot a(s) \quad s \in G\,.$$

The stated result thus follows from Theorem 1.2 and the straightforward identification $\mathrm{sp}(L) = \{\ell(s) \in [0, +\infty) : s \in \Gamma\}$.

□

## References


[1] M. Bozejko, T. Januszkiewicz, R.J. Spatzier, *inifinite Coxeter groups do not have the Kazhdan's property*, J. Operator Theory, **19**(1), (1988), 63-67.

[2] P.-A. Cherix, M. Cowling, P. Jolissaint, P. Julg, A. Valette, *Groups with the Haagerup Property. Gromov a-T-menability* Progress in Mathematics **197**, Birkhauser Verlag, Basel - Boston - Berli, 2001.

[3] F. Cipriani, *Dirichlet forms in noncommutative spaces. Quantum Potential Theory*, Lect. Notes in Math., **1954**, 161-276, Springer, Berlin, 2008.

[4] F. Cipriani, U. Franz, A: Kula, *Symmetries of Lévy processes on compact quantum groups, their markov semigroups and potential theory*, J. Funct. Anal., **266**, (2014), 2789-2844.

[5] F. Cipriani, J.-L. Sauvageot, *Derivations as square roots of Dirichlet forms*, J. Funct. Anal., **201**, (2003), 78-120.

[6] F. Cipriani, J.-L. Sauvageot, *variations in Noncommutative Potential Theory: finite energy states, potentials and multipliers*, Trans. Amer. Math. Soc., **367**(7), (2015), 48374871.

[7] F. Cipriani, J.-L. Sauvageot, *Amenability and sub-exponential spectral growth rate of Dirichlet forms on von Neumann algebras*, in preparation.

[8] J. Dixmier, *Operateur de rang fini dans les representation unitaires*, Publ. Math. I.H.E.S., **6**, (1960), 13-35.

[9] M. Gromov, *Groups of polynomial growth and expanding maps*, Publ. Math. I.H.E.S., **53**, (1981), 53-73.

[10] M. Gromov, *Rigid transformations groups*, In Géométrie différentielle, variétés complexes, felleuitage riemanniens (Paris 1986), Travaux en cours **33**, (1988), 65-139, D. Bernard and Y Choquet-Bruhat , editors, Hermann Paris (1988).

[11] U Haagerup, *An example on nonnuclear $C^*$-algebra, which has the metric approximation property*, Invent. Math., **53**(3), (1979), 279-293.

[12] P. de la Harpe, *Topics in Geometric group theory* Chicago Lectures in Mathematics, The University of Chicago Press, Chicago and London, 2000.

[13] V. Losert, *On the structure of groups with polynomial growth*, Math. Z., **195**, (1987), 109-117.



Dipartimento di Matematica, Politecnico Milano Piazza Leonardo da Vinci 33 20132 Milano Italy

*E-mail address*: fabio.cipriani@polimi.it

Institut de Mathématiques, CNRS Université Denis Diderot F-75205 Paris Cedex 13, France

*E-mail address*: jlsauva@math.jussieu.fr